


To appear in *Dwarf Galaxies and Cosmology*, the Proceedings of the XVIIIth Moriond Astrophysics Meeting, Les Arcs, Savoie, France, 14th–21st March, 1998, eds. T. X. Thuân, C. Balkowski, V. Cayatte and J. Trân Thanh Vân, publ. Editions Frontières, Gif-sur-Yvette.

DWARF SPHEROIDAL GALAXIES IN THE VIRGO CLUSTER

J. B. Jones ¹, S. Phillipps ¹, J. M. Schwartzberg ¹, Q. A. Parker ²

¹ *Astrophysics Group, Department of Physics, University of Bristol,
Tyndall Avenue, Bristol, BS8 1TL, United Kingdom.*

² *Anglo-Australian Observatory, Siding Spring, Coonabarabran,
New South Wales 2357, Australia.*



Abstract

We present a study of the smallest and faintest galaxies found in a very deep photographic R band survey of two regions of the Virgo Cluster, totalling 3.2 square degrees, made with the UK Schmidt Telescope. The objects we detect have the same physical sizes and surface brightnesses as Local Group dwarf spheroidal galaxies. The luminosity function of these extremely low luminosity galaxies (down to $M_R \simeq -11$ or about $5 \times 10^{-5} L_\star$) is very steep, with a power law slope $\alpha = -2.2$.

1 Introduction

The properties of the faint end of the luminosity function are poorly constrained outside the Local Group, both in terms of the numbers and characteristics of the galaxies. Some studies have found evidence for very steep luminosity functions in both cluster [1][2] and field [3] environments. The importance of very low luminosity galaxies in understanding galaxy formation and evolution [2] demands that progress is made in identifying and studying low luminosity dwarfs in new environments [4].

This survey extends the study of extremely low luminosity galaxies to the Virgo Cluster (to $M_R = -11$ mag), into the regime of Local Group dwarf spheroidals. It extends the definitive study of the Virgo Cluster by Binggeli, Sandage & Tammann [5] by 3 mag beyond their completeness limits and complements the survey of large, low surface brightness galaxies of Impey, Bothun & Malin [6]. By coadding multiple photographic exposures with the UK Schmidt Telescope, galaxies are detected having central surface brightnesses as faint as 25 R mag arcsec⁻², equivalent to 26.5 B mag arcsec⁻². We report here results from an initial survey [7] covering 3.2 deg².

2 The Data and Image Detection

The data used here [7][8] are part of a larger photographic survey [9] of a ten-degree square region of the Virgo Cluster using the extremely fine grained, highly efficient Tech Pan film on the 1.2m UK Schmidt Telescope [10]. Six individual long (1 to 1.5 hour) exposures of the same area, the South-East quadrant of the Virgo Cluster, were scanned with the SuperCOSMOS automatic measuring machine at the Royal Observatory Edinburgh. Two $1.3^\circ \times 1.3^\circ$ subregions were used for the initial survey, one near the cluster centre, one further out. The digitised data from the six separate films were sky subtracted and were matched in intensity [11] before coaddition using median stacking. This provided an equivalent exposure time of about 7 hours, while the coadded data are free of artefacts such as emulsion defects, dust particle images and satellite trails that affect only one film.

Image detection was performed with a connected-pixel algorithm using a detection threshold $\mu_{lim} = 25.5 \text{ mag arcsec}^{-2}$ and a minimum area above the threshold of $A_{lim} = 11$ pixels. Each image thus had a minimum S/N ratio of 10 and has a magnitude $R \leq 22$. This detection limit is $0.3 \text{ mag arcsec}^{-2}$ fainter than previous surveys. Around 28 000 images were detected in both fields and the large minimum area ensured that few were spurious. This was confirmed by comparison of a small subregion of the photographic field with a CCD image taken on the Anglo-Australian Telescope: 160 of 162 images visible on the film were matched on the CCD frame, including all those which contributed to determining the luminosity function in Section 4. Scale lengths and surface brightnesses have been measured for each image from the isophotal signals and areas at the detection threshold, assuming exponential light profiles.

3 Identifying Virgo dwarfs

Virgo Cluster dwarf elliptical and dwarf spheroidal galaxies can be characterised by their image sizes and surface brightnesses: they are larger in general than the majority of background images at a given central surface brightness. As illustrated in Figure 1, it is possible to use the position of images in the magnitude – surface brightness plane to identify a sample rich in Virgo dwarfs. Figure 1a compares the distribution of images in a Virgo field with that of the background population. The background data are based on AAT CCD observations of a field around the South Galactic Pole. The background population dominates strongly at scale lengths $\alpha \lesssim 1.8 \text{ arcsec}$. In contrast, a majority of galaxies having scale lengths $\alpha > 2 \text{ arcsec}$ and central surface brightnesses $\mu_0 > 22.0 \text{ mag arcsec}^{-2}$ are expected to be Virgo dwarfs.

The properties of the observed galaxies can be compared with those of Local Group objects. Figure 1b shows the positions that Local Group dSphs [12] would occupy in the magnitude – surface brightness diagram were they located at the distance of the Virgo Cluster: many of the local dSphs could be distinguished unambiguously from the background population. For comparison, the regions occupied by the galaxies in the samples of Binggeli, Sandage & Tammann [5] and of Impey, Bothun & Malin [6] are shown in the figure; neither of these other studies reaches the small, low surface brightness galaxies found here.

4 The luminosity Function

Information about number densities of genuine Virgo Cluster galaxies of various types can be derived by subtracting the background contribution from the total number of observed galaxies at each point in the magnitude – surface brightness plane. This technique has been used to




Figure 1: (a) The magnitude – central surface brightness plane for galaxies detected in the Virgo Schmidt survey compared with the SGP CCD field used for background subtraction. For clarity, galaxies having scale lengths $\alpha < 2$ arcsec (where the background population dominates) have not been plotted. Loci are shown corresponding to exponential galaxies having scale lengths $\alpha = 1, 2, 10$ and 30 arcsec. The selection boundary imposed on the Schmidt data by the detection methods is shown as the dashed curve. The bold lines show the limits of the sample used to determine the luminosity function in Section 4. (b) As Figure 1a, but showing the properties Local Group dSphs would have if seen at the distance of the Virgo Cluster. The solid lines mark the regions of the plane covered by the Binggeli *et al.* [5] and Impey *et al.* [6] samples. Throughout this paper we adopt a distance modulus of 31.3 mag for the cluster.

determine the luminosity function for a subsample of Virgo dwarfs, defined by $\alpha \geq 3.0$ arcsec and central surface brightnesses $22.0 \text{ mag arcsec}^{-2} \leq \mu_0 \leq 24.5 \text{ mag arcsec}^{-2}$, in each of the two fields studied. This scale length limit ensures that the background galaxy contamination is minimal (about 4%), reducing difficulties which might be caused by comparing data having different point-spread functions. These sample limits are shown in Figure 1a.

The luminosity functions [7] for the two fields are presented in Figure 2. The inner field is centred close to M87 and samples the population in the core of the cluster. The area immediately around M87, having a higher effective background light level, has been excluded from the analysis; the area surveyed is 1.58 deg^2 . The outer cluster field lies 3.1 deg to the south-southeast of M87 and is 1.61 deg^2 in area. The inner and outer cluster fields have very similar luminosity functions. Both have steep faint-end slopes: the combined data set has $\alpha = -2.26 \pm 0.13$ over the magnitude range $15.5 < R < 20.0$. The downturn at the last low luminosity point is a consequence of the sample limits (defined by scale length and surface brightness, not by magnitude). At the bright end, the luminosity functions are consistent with that of Binggeli, Sandage & Tammann [13][14] (although a comparison is complicated by the slightly different samples and filters used in the two surveys). The density of galaxies is marginally greater in the outer cluster field than in the core (by a factor of 30%), which is consistent with a depletion of dwarfs in the core, as has been observed in the Coma Cluster [15].

The survey is currently being extended to the full ten-degree square region centred on

the cluster core. Photographic data in an off-cluster field will provide improved background subtraction, allowing a luminosity function to be constructed which extends further into the region of Figure 1 occupied by field galaxies than has been possible here.




Figure 2: The absolute magnitude distribution of the sample of Virgo dwarfs of Section 4 for the (a) outer and (b) inner fields. Error bars shown are based on Poissonian statistics. The luminosity function of Sandage, Binggeli & Tammann [13][14] is also shown (open triangles) for the Virgo Cluster Catalog galaxies which overlap the inner cluster field (assuming $B-R = 1.5$).

References

- [1] Smith R.M., Driver S.P., Phillipps S., 1997, *MNRAS* **287**, 415
- [2] Trentham N., 1998, this volume (and astro-ph/9804013).
- [3] Loveday J., 1997, *Astrophys. J.* **489**, 29
- [4] Caldwell N., Armandroff T.E., Da Costa G.S., Seitzer P., 1998, *Astron. J.* **115**, 535
- [5] Binggeli B., Sandage A., Tammann G.A., 1985, *Astron. J.* **90**, 1681
- [6] Impey C., Bothun G., Malin D., 1988, *Astrophys. J.* **330**, 634
- [7] Phillipps S., Parker Q.A., Schwartzberg J.M., Jones J.B., 1998, *Astrophys. J.* **493**, L59
- [8] Schwartzberg J.M., 1996, Ph.D. Thesis, University of Bristol
- [9] Schwartzberg J.M., Phillipps S., Parker Q.A., 1995, *Astr. Astrophys.* **293**, 332
- [10] Phillipps S., Parker Q.A., 1993, *MNRAS* **265**, 385
- [11] Schwartzberg J.M., Phillipps S., Parker Q.A., 1996, *Astr. Astrophys. Suppl. Ser.* **117**, 179
- [12] Caldwell N., Armandroff T.E., Seitzer P., Da Costa G.S., 1992, *Astron. J.* **103**, 840
- [13] Sandage A., Binggeli B., Tammann G.A., 1985, *Astron. J.* **90**, 1759
- [14] Binggeli B., Sandage A., Tammann G.A., 1988, *Ann. Rev. Astron. Astrophys.* **26** 509
- [15] Thompson L.A., Gregory S.A., 1993, *Astron. J.* **106**, 2197